



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

$a, a+x, a+2x, a+3x, \dots, a+(n-1)x$ , is an arithmetical progression of  $n$  terms, their sum of squares is  $a^2 + (a+x)^2 + (a+2x)^2 + (a+3x)^2 + \dots$

$+ [a+(n-1)x]^2 = S = na^2 + (n^2-n)ax + \frac{1}{6}(2n^3-3n^2+n)x^2$  by method of difference. This sum is made a rational square by the method given in Encyclopedia Britannica, Vol. I, Algebra article 121, 9th edition. When  $n$  is a square number  $na^2 + (n^2-n)ax + \frac{1}{6}(2n^3-3n^2+n)x^2 = (n^{\frac{1}{2}}a + hx)^2$

$= na^2 + 2ahnx + \frac{1}{6}(2n^3-3n^2+n)x^2$ . Then  $x = \frac{2ahn^{\frac{1}{2}} - (n^2-n)a}{\frac{1}{6}(2n^3-3n^2+n) - h^2} = \frac{2}{3}$  for  $a=1$ ,

$n=9, h=14$ . The algebraic value of  $x$  in expressions for  $S_1$  after clearing denominatives gives a general solution, the numerical  $\frac{2}{3}$ , gives

$$2^2 + 5^2 + 8^2 + 11^2 + 14^2 + 17^2 + 20^2 + 23^2 + 26^2 = 48^2.$$

Also solved by F. P. Matz, G. B. M. Zerr, J. H. Drummond, C. D. Schmitt, J. Scheffer, and M. A. Gruber

---

## PROBLEMS.

---

30. Proposed by COOPER D. SCHMITT, Knoxville, Tennessee.

$A$  and  $B$  are two integers,  $A$  consisting of  $2m$  figures each being 1, and  $B$  consisting of  $m$  figures each being 4. Prove that  $A + B + 1$  is a square.

31. Proposed by M. A. GRUBER, War Department, Washington, D. C.

How many scalene triangles, of integral sides, can be formed with an altitude of 12? How many isosceles triangles?

---

## AVERAGE AND PROBABILITY.

---

Conducted by B.F.FINKEL, Kidder, Missouri. All contributions to this department should be sent to him.

---

## SOLUTIONS OF PROBLEMS.

---

17. Proposed by A. L. FOOTE, No. 80, Broad St. New York.

A person 30 years of age has an annuity for 10 years, the present worth of which is \$1000, provided he lives but ten years; for, if he dies, the annuity ceases. What is the annuity worth, on the assumption that 75 out of every 4385 persons die annually, between the ages 30 and 40 years?

Solution by B. F. FINKEL, A. M., Professor of Mathematics and Physics, Kidder Institute, Kidder, Missouri.

Let  $S$  = the annuity, the present value,  $P$ , of which, for 10, =  $n$ , years is \$100.

Then  $P = \frac{S}{R-1} \left( \frac{R^n-1}{R^n} \right)$ , whence  $S = \frac{rPR^n}{R^n-1}$ , where  $R=1+r$ . If

we assume  $r=.0$ , we find  $S=\$149.0294$  nearly.

The limit of  $B$ 's life is  $4385 \div 75 = 58\frac{1}{3}$  years,  $=l$ .  $\therefore$  the probability that  $B$  will be living at the end of 1 year is  $\frac{l-1}{l}$ ; at the end of two years,  $\frac{l-2}{l}$ ; at the end of three years,  $\frac{l-3}{l}$ ; etc.

The present worth of  $S$ ,  $=\$149.20207$ , due in 1, 2, 3, 4, etc., years is  $\frac{S}{R}$ ,  $\frac{S}{R^2}$ ,  $\frac{S}{R^3}$ , etc., to  $\frac{S}{R^n}$ . The present worth of  $S$  due at the end of any year multiplied by the probability of  $B$ 's living to the end of that year is the actual value of  $S$ .

$$\therefore S' = \frac{S(l-1)}{Rl} + \frac{S(l-2)}{lR^2} + \frac{S(l-3)}{lR^3} + \dots + \frac{S(l-n)}{lR^n} \dots (1).$$

$$\frac{S'}{R} = \frac{S(l-1)}{lR^2} + \frac{S(l-2)}{lR^3} + \frac{S(l-3)}{lR^4} + \dots + \frac{S(l-n)}{lR^{n+1}} \dots (2), \text{ by multiplying (1) by } \frac{1}{R}.$$

$$\begin{aligned} \therefore S' \left( \frac{R-1}{R} \right) &= \frac{S(l-1)}{Rl} - \frac{S}{lR^2} - \frac{S}{lR^3} - \dots - \frac{S(l-n)}{lR^{n+1}}, \\ &= \frac{S}{l} \left\{ \frac{l}{R} - \frac{1}{R} - \frac{1}{R^2} - \frac{1}{R^3} \dots - \frac{1}{R^n} - \frac{l-n}{R^{n+1}} \right\}, \\ &= \frac{S}{l} \left\{ \frac{l}{R} - \frac{l-n}{R^{n+1}} - \frac{1}{R} - \frac{1}{R^2} \dots - \frac{1}{R^n} \right\}, \\ &= \frac{S}{l} \left\{ \frac{l}{R} - \frac{l-n}{R^{n+1}} - \frac{1}{R} \left[ 1 - \frac{1}{R^n} \right] \div \frac{R-1}{R} \right\}, \\ &= \frac{S}{R} \left\{ \left( 1 - \frac{l-n}{lR^n} \right) - R \left( 1 - \frac{1}{R^n} \right) \div (R-1)l \right\}, \text{ whence } S' = S \div r \left\{ \left[ 1 - (l-n) \div \right. \right. \\ &\left. \left. lR^n \right] - R \left( 1 - \frac{1}{R^n} \right) \div rl \right\}, = \$911.881029. \end{aligned}$$

No solutions of this problem were received from our contributors.

#### 18. Proposed by H. W. DRAUGHON, Clinton, Louisiana.

The probability that  $A$  will speak the truth is twice the probability that  $B$  will, in an independent statement, speak the truth; but, if  $A$  exerts his influence, the probability is that  $B$  will agree with him in any statement. What is the probability of the truth of their concurrent testimony, the chances being equal that  $A$  may or may not be interested in the matter?

Solution by P. H. PHILBRICK, C. E., Lake Charles, Louisiana.

1. Suppose that  $A$  is not interested in the matter. Let  $x$  = the probability of the truth of any one of  $B$ 's statements. Then  $2x$  = the probability of the truth of any one of  $A$ 's statements. The event did occur if both witnesses tell the truth the probability of which is  $x \times 2x = 2x^2$ .